

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
8 March 2001 (08.03.2001)

PCT

(10) International Publication Number  
**WO 01/15542 A1**

- (51) International Patent Classification<sup>7</sup>: A23D 7/00 Bunka, Sumida-ku, Tokyo 131-8501 (JP). YASUNAGA, Koichi [JP/JP]; Kao Corporation, Research Laboratories, 2-1-3, Bunka, Sumida-ku, Tokyo 131-8501 (JP).
- (21) International Application Number: PCT/JP00/05583
- (22) International Filing Date: 21 August 2000 (21.08.2000) (74) Agents: FURUYA, Kaoru et al.; Hamacho-Hanacho Building 6th Floor, 2-17-8, Nihonbashi-Hamacho, Chuo-ku, Tokyo 103-0007 (JP).
- (25) Filing Language: English
- (26) Publication Language: English (81) Designated States (*national*): BR, CA, CN, US.
- (30) Priority Data: 11/243432 30 August 1999 (30.08.1999) JP (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (71) Applicant (*for all designated States except US*): KAO CORPORATION [JP/JP]; 14-10, Nihonbashi-Kayabacho 1-chome, Chuo-ku, Tokyo 103-0025 (JP). Published:  
— With international search report.
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): MASUI, Kenji [JP/JP]; Kao Corporation, Research Laboratories, 2-1-3, For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: WATER-IN-OIL TYPE EMULSIFIED FAT AND/OR OIL COMPOSITION



(57) Abstract: The present invention provides a water-in-oil type emulsified fat and/or oil composition of which emulsification is stable in spite of containing a high content of water and which is excellent during storage in physical properties and feeling on eating. That is, the present invention provides a water-in-oil type emulsified oil and/or fat composition which is composed of (1) the oil phase comprising 35 to 95 % by weight, based on the total oils and fats, of a diglyceride(s) having its increasing melting point of lower than 20 °C and the balance of triglycerides having fatty acid parts comprising 13 to 60 % by weight of palmitic acid and 5 % by weight or less of fatty acids having 12 or less carbon atoms, polymorph of the triglycerides being stable in the form of  $\beta'$  and (2) the aqueous phase based on water.

WO 01/15542 A1

## DESCRIPTION

## WATER-IN-OIL TYPE EMULSIFIED FAT AND/OR OIL COMPOSITION

## Technical Field

The present invention relates to a water-in-oil type emulsified fat and/or oil composition which is utilized preferably and suitably in a margarine, a fat spread, a shortening etc.

## Background Art

As people are health-oriented in recent years, a product being low-calorie- and low-oil-content- type is desired for a margarine or a fat spread.

In general, the method for lowering an oil content includes raising the ratio of an aqueous phase to provide a water-in-oil type emulsified fat and/or oil composition which is a high content of water-containing type. However, when usual fats and oils of triglycerides are used in the oil phase, the emulsified state becomes unstable with an increasing ratio of the aqueous phase. Thus, in during the production step, the aqueous phase tends to separate from the oil phase in a mixing tank for the aqueous at the step of preliminary emulsification and oil phases and the phase reversal from W/O to O/W tends to occur. Further, oozing of water and oil-off is caused during storage. When bulk-handling is carried out, the emulsification is destroyed, and separation, decay or the like of the aqueous is carried out thereby. Therefore, a product exhibiting

a satisfied performance could not be obtained.

To solve these disadvantages, methods for using various emulsifiers such as a sucrose fatty acid ester (JP-B 56-10014), a glycerol fatty acid ester (US-A 3889005) and a polyglycerol fatty acid ester (JP-A 58-170432) have been proposed.

However, when the above-mentioned emulsifier is used, although the water-in-oil type emulsified fat and/or oil composition which is a high content of water-containing type can be produced, the obtained product is badly stable by the day during storage so that water oozes and the emulsification is destroyed upon spreading, therefore the following separation of the aqueous phase is generated.

Meanwhile, various methods for blending a large amount of diglycerides (glycerol di-fatty-acid esters) with the oil phase to solve the above-described disadvantages have also been proposed in, for example, JP-A 63-301743, JP-A 61-63242, WO 95/22257, WO 96/32022 and US-A 5879735.

Although a stable water-in-oil type emulsified fat and/or oil composition can be produced by these methods, the product is worsened in viewpoint of the physical property during storage. Then, the product is made to be crumbly and so on, it is poor in feeling on eating, and its spreadability is not satisfied.

#### Disclosure of the Invention

The present inventors developed a water-in-oil type emulsified fat and/or oil composition of which emulsification is stable in spite of containing a high content of water and which is

excellent in physical properties and feeling on eating during storage.

The present invention provides an water-in-oil type emulsified fat and/or oil composition which is composed of (1) the oil phase comprising 35 to 95 % by weight, based on the total oils and fats, of a diglyceride(s) having its increasing melting point of lower than 20 °C and the balance of triglycerides having fatty acid parts comprising 13 to 60 % by weight of palmitic acid and 5 % by weight or less of fatty acids having 12 or less carbon atoms, polymorph of the triglycerides being stable in the form of  $\beta'$  and (2) the aqueous phase based on water. Thus, the above-mentioned problem of the prior art has been solved.

#### Brief Description of the Drawing

Fig. 1 shows a triangular chart exhibiting a ratio of palmitic acid, stearic acid and an unsaturated fatty acid having 16 or more carbon atoms among fatty acids constituting triglycerides in the oil phase in the water-in-oil type emulsified fat and/or oil composition obtained in the Examples, wherein ① to ④ show the results in Examples 1 to 4 and ①' to ③' show the results in Comparative Examples 1 to 3.

#### Modes for Carrying Out the Invention

The fatty acid constituting the diglyceride having its increasing melting point of lower than 20 °C (preferably -20 to 15 °C) is preferably an unsaturated fatty acid having 16 to 22 carbon

atoms, and its content is 70 % by weight or more, more preferably 80 % by weight or more, based on the fatty acid constituting the diglyceride.

As the starting fat or oil used in the diglyceride, a fat or oil containing a large amount of an unsaturated fatty acid-residue, for example, including a vegetable (or plant) fat or oil such as safflower oil, olive oil, cottonseed oil, rapeseed oil, corn oil, soybean oil and palm oil; further, an animal fat or oil such as lard, beef-tallow and butter; and a fractional oil thereof, a transesterified oil thereof and a hydrogenated oil (hardened oil) thereof may be used. The diglyceride is obtained by ester-exchanging reaction (, ester-interchanging reaction or transesterification) of a mixture of one or more of these fats and oils and glycerol in the presence of a catalyst or by ester-exchanging reaction of a mixture of a fatty acid composition having the high levels of unsaturated fatty acids and glycerol in the presence of a catalyst.

An excess of monoglycerides formed in a mixture of the generated diglycerides can be removed by a molecular distillation method. The monoglycerides are desirably removed to make it as small as possible. It is more preferable that the content thereof is 5 % by weight or less, preferably 2 % by weight or less, .

From the viewpoint of the stability in the emulsification, the diglycerides are blended in an amount of 35 to 95 % by weight and preferably 45 to 90 % by weight based on the total fats and oils.

Then, the starting materials of triglycerides whose

crystalline polymorph is stable in the form of  $\beta'$  in the present invention are edible fats and/or oils, and there is no particular limitation insofar as the content of a fatty acid having 12 or less carbon atoms is 5 % by weight or less based on the total fatty acids for constituent. Namely, there may be used one or more selected from vegetable fats and oils including soybean oil, rapeseed oil, palm oil, corn oil, cottonseed oil, coconut oil and palm kernel oil; and animal fats and oils including lard, fish oil and milk fat; and those obtained by hydrogenation or ester-exchange of these oils. When the content of a fatty acid having 12 or less carbon atoms is more than 5 % by weight in the fatty acids constituting triglycerides, a uniquely soapy smell is easily generated by the low-temperature deterioration and hydrolysis of an lauric fat and/or oil to cause particularly the flavor deterioration in margarine, spread etc. during storage.

In the present invention, the term  $\beta'$  in the crystalline polymorph of a fat and/or oil indicates a general tendency for a majority of fats and oils to be crystallized, stable and continued in the  $\beta'$  phase and doesn't indicate the crystalline polymorph of the fat and/or oil not being able to transfer from the  $\beta'$  phase to  $\beta$  phase under a proper condition. The type of crystalline polymorph of fats and oils can be analyzed and identified by X-ray diffraction pattern thereof, and a method thereof is described in general literatures (e.g., "Science of edible fat and oil", p. 201, published by Saiwai Shobo).

In the water-in-oil emulsified fat and/or oil composition of

the present invention, the content of palmitic acid in fatty acids constituting triglycerides is desirably 13 to 60 % by weight so that the crystalline polymorph of triglycerides in the oil phase containing diglycerides is stable in the form of  $\beta'$ . The remained fatty acids are myristic acid and a saturated or unsaturated fatty acid having 16 to 22 carbon atoms. Further, it is more preferable that a ratio of palmitic acid, stearic acid and an unsaturated fatty acid having 16 or more carbon atoms among fatty acids constituting triglycerides is in the range surrounded by points A (13, 2, 85), B (13, 57, 30), C (60, 10, 30) and D (60, 2, 38) in the triangular chart shown in Fig. 1, whereby it is recognized that the melting behavior in the mouth is significantly improved and the hydration and whipping are also improved. It is particularly preferable that the ratio is in the range surrounded by points A (13, 2, 85), E (13, 47, 40), F (50, 10, 40) and G (50, 2, 48).

The water-in-oil type emulsified fat and/or oil composition of the present invention may be prepared in a usual manner, and the ratio of the aqueous phase based on water to the oil phase by weight can be made within the range of the oil phase : the aqueous phase = 20 : 80 to 85 : 15 (preferably 40 : 60 to 85 : 15).

A dairy product, an emulsifier, a flavor ingredient or the like as a sub-component in the oil phase may be blended with the composition of the present invention. Further, a flavor ingredient, milk protein, starch, a thickening polysaccharide, a thermoplastic protein (gelatin etc.) or the like may be blended with the aqueous phase based on water.

Further, an anti-oxidant including tocopherol, an ascorbyl ester such as vitamin C palmitate and vitamin C stearate, a tea leaf, a herb such as rosemary, a natural anti-oxidizing component extracted from a leaf or a root of a peach may be used together in order to inhibit the deterioration of the water-in-oil type emulsified fat and/or oil composition.

#### Examples

Examples 1 to 4 and Comparative Examples 1 to 3.

#### [Preparation of diglyceride]

While the fatty acid obtained from a soybean oil and glycerol were mixed at the molar ratio of 2 : 1, they were reacted at 70 °C for 3 hours in the presence of a catalyst as a commercial lipase preparation of immobilized 1,3-position-selective lipase (Lipozyme 3A™ provided by Novo Industry A. S.). During the reaction, the inside pressure of the system was reduced to 0.26 kPa in order to remove water formed by esterification. After the lipase preparation was removed by filtration from the reaction product, the filtrate was subjected to molecular distillation, decoloration and deodorization to obtain a diglyceride composition (consisting of diglyceride content of 88 % by weight, monoglyceride content of 1.3 % by weight and triglyceride content of 10.7 % by weight).

The above-mentioned composition was fractionated by column chromatography (a triglyceride fraction was eluted with hexane from a column packed with Wako Gel C-200 provided by Wako Pure Chemicals, Industries and then a diglyceride fraction was obtained with

hexane/ether). As a result of measurement of the melting point of the diglyceride fraction, the open tube melting point was 5 °C.

According to the formulation in Table 1, the oil phase and the aqueous phase were mixed and emulsified, and the obtained emulsion was rapidly cooled and plasticized in a usual manner to obtain the water-in-oil type emulsified oil and/or fat compositions in Examples 1 to 4 and Comparative Examples 1 to 2.

In the obtained water-in-oil type emulsified oil and/or fat composition, the content of diglycerides and monoglycerides based on the total fats and oils, as well as the composition of fatty acids in the triglyceride, are shown in Table 2.

The above-mentioned compositions were stored at 5 °C for 30 days and then evaluated for crystalline polymorph of fats and oils, the stability in the emulsification (the state of texture and separation in the aqueous phase at the time for spread), spreadability (coating properties) and feeling on eating (smoothness). The results are shown in Table 1.

The evaluation criteria are as follows.

- Crystalline polymorph of fat(s) and/or oil(s): X-ray diffraction was carried out in a usual manner.
- Stability in the emulsification: The produced emulsified fat and/or oil composition was coated on bread etc. and then evaluated for separation of the aqueous phase under the following criteria.

○: Water was not separated.

△: Water was slightly separated to ooze.

X: Water was separated.

• Spreadability: The produced emulsified fat and/or oil composition was coated on bread etc. and then evaluated for performance (degree of easiness to spread, uniform texture and smoothness) under the following criteria.

○: Spread thin and continuously.

△: Hardly spread thin and continuously.

×: Not coated thin and continuously.

• Feeling on eating: The feeling (smoothness) upon eating of the produced emulsified fat and/or oil composition was evaluated as organoleptic property under the following criteria.

○: Smooth and not crumbly in the mouth.

△: Slightly crumbly and slightly rough upon eating.

×: Evidently crumbly and rough upon eating.

Table 1

	Example 1	Example 2	Example 3	Example 4	Comparative Example 1	Comparative Example 2	Comparative Example 3
Oil phase (% by weight)							
Diglyceride composition	31.9	31.9	48.6	33.9	31.9	31.9	31.9
Soybean oil							
Palm hydrogenated (or hardened) oil (IV 20)	1.6	4.8	18.7				
Palm hydrogenated (or hardened) oil (IV 2)			2.1	4			
Soybean hydrogenated (or hardened) oil (IV 43)	6.4	3.2		2	8	8	8
Soybean hydrogenated (or hardened) oil (IV 1.3)							
Stearic acid monoglyceride			0.5				
Vitamin E			0.02				
Carotene			0.002				
Butter flavor	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Aqueous phase (% by weight)							
Distilled water	60	60	28.4	60	60	60	60
Skim milk powder			0.3				
Common salt			1.3				
Crystal form of fat or oil	$\beta'$	$\beta'$	$\beta'$	$\beta'$	$\beta$	$\beta$	$\beta$
Stability of emulsification	O	O	O	O	X	O	O
Spreadability	O	O	O	O	X	$\Delta$	$\Delta$
Feeling on eating	O	O	O	O	X	$\Delta$	X

Table 2


	Example 1	Example 2	Example 3	Example 4	Comparative 1	Comparative 2	Comparative 3
Diglyceride content (% by weight)	70.4	70.3	61.6	74.8	2.0	70.4	70.4
Monoglyceride content (% by weight)	1.0	1.1	1.6	1.1	0.0	1.0	1.0
Fatty acid composition in the triglyceride fraction (% by weight)							
C14	0.2	0.5	1.0	0.6	0.0	0.1	0.1
C16	15.9	25.2	37.7	24.7	11.1	11.3	11.2
C18	10.6	9.0	11.4	27.3	6.3	11.4	62.8
C18:1	54.6	46.7	37.2	24.6	34.5	58.5	7.9
C18:2	16.2	16.2	10.8	19.9	42.1	16.1	15.8
C18:3	2.5	2.5	1.8	2.8	6.0	2.5	2.2

It is evident from Table 1 that the emulsified fat and/or oil composition of the present invention exhibits stable emulsification and is excellent in stability of physical property during storage.

## CLAIMS

- 1, An water-in-oil type emulsified fat and/or oil composition which is composed of (1) the oil phase comprising 35 to 95 % by weight, based on the total oils and fats, of a diglyceride(s) having its increasing melting point of lower than 20 °C and the balance of triglycerides having fatty acid parts comprising 13 to 60 % by weight of palmitic acid and 5 % by weight or less of fatty acids having 12 or less carbon atoms, polymorph of the triglycerides being stable in the form of  $\beta'$  and (2) the aqueous phase based on water.
- 2, The water-in-oil type emulsified fat and/or oil composition as claimed in Claim 1, wherein a ratio of palmitic acid, stearic acid and an unsaturated fatty acid having 16 or more carbon atoms among fatty acids constituting triglycerides is in the range surrounded by points A (13, 2, 85), B (13, 57, 30), C (60, 10, 30) and D (60, 2, 38) in the triangular chart in Fig. 1.

Fig. 1



An unsaturated fatty acid  
having 16 or more carbon atoms

# INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/JP 00/05583

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A23D7/00

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A23D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, FSTA

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 32022 A (UNILEVER NV ;UNILEVER PLC (GB)) 17 October 1996 (1996-10-17) cited in the application example 2 claims 1-7	1
Y	EP 0 425 958 A (KAO CORP) 8 May 1991 (1991-05-08) page 5, line 35 - line 49 page 7, line 39 -page 9, line 35	1
Y	US 5 684 230 A (FEHR WALTER R ET AL) 4 November 1997 (1997-11-04) column 1, line 34 -column 2, line 6 claims 1,2	1
	--- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

**\* Special categories of cited documents:**

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*8\* document member of the same patent family

Date of the actual completion of the international search

20 November 2000

Date of mailing of the international search report

27/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Dekeirel, M

# INTERNATIONAL SEARCH REPORT

Intern. Appl. Application No

PCT/JP 00/05583

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>EP 0 430 329 A (UNILEVER NV ;UNILEVER PLC (GB)) 5 June 1991 (1991-06-05)  page 2, line 31 - line 36  page 2, line 25 - line 29</p>	1
A	<p>WO 98 34494 A (BEYER RICHARD ;UNIV OTAGO (NZ); KIWITECH LIMITED (NZ); SILCOCK PAT) 13 August 1998 (1998-08-13)  page 6, line 27 -page 7, line 2  page 10, line 13 - line 18  example 5  claims 1,7,8,44,45</p>	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/JP 00/05583

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9632022 A	17-10-1996	AU 5496896 A	30-10-1996
EP 0425958 A	08-05-1991	DE 69012737 D	27-10-1994
		ES 2059950 T	16-11-1994
		JP 2862981 B	03-03-1999
		JP 3236743 A	22-10-1991
		US 5077077 A	31-12-1991
US 5684230 A	04-11-1997	US 5602311 A	11-02-1997
		US 5663485 A	02-09-1997
		US 5714672 A	03-02-1998
		US 5684231 A	04-11-1997
		US 5850029 A	15-12-1998
		US 5557037 A	17-09-1996
		US 5795969 A	18-08-1998
EP 0430329 A	05-06-1991	EP 0398411 A	22-11-1990
		AT 117512 T	15-02-1995
		AU 636414 B	29-04-1993
		AU 6677790 A	30-05-1991
		CA 2030508 A,C	23-05-1991
		DE 69016368 D	09-03-1995
		DE 69016368 T	14-06-1995
		DK 430329 T	19-06-1995
		FI 905696 A	23-05-1991
		IE 67654 B	17-04-1996
		JP 3272645 A	04-12-1991
		JP 7089869 B	04-10-1995
		NO 905048 A	23-05-1991
		TR 26084 A	15-12-1994
		US 5244688 A	14-09-1993
		ZA 9009383 A	29-07-1992
		AT 83126 T	15-12-1992
		AT 96614 T	15-11-1993
		CA 2025539 A,C	30-03-1991
		DE 69004396 D	09-12-1993
		DE 69004396 T	10-03-1994
WO 9834494 A	13-08-1998	AU 6007498 A	26-08-1998